

An Efficient and Rapid Synthesis of 1,4-Dihydropyrano[2,3-c]Pyran and 1,4-Dihydropyrano[2,3-c]Quinoline Derivatives Using Copper Nanoparticles Grafted on Carbon Microspheres

Nitishkumar S. Kaminwar, Sunil U. Tekale, Rajkumar U. Pokalwar, László Kótai & Rajendra P. Pawar

To cite this article: Nitishkumar S. Kaminwar, Sunil U. Tekale, Rajkumar U. Pokalwar, László Kótai & Rajendra P. Pawar (2021): An Efficient and Rapid Synthesis of 1,4-Dihydropyrano[2,3-c]Pyran and 1,4-Dihydropyrano[2,3-c]Quinoline Derivatives Using Copper Nanoparticles Grafted on Carbon Microspheres, *Polycyclic Aromatic Compounds*, DOI: [10.1080/10406638.2021.1950194](https://doi.org/10.1080/10406638.2021.1950194)

To link to this article: <https://doi.org/10.1080/10406638.2021.1950194>



Published online: 21 Jul 2021.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)



An Efficient and Rapid Synthesis of 1,4-Dihydropyrano [2,3-c]Pyran and 1,4-Dihydropyrano[2,3-c]Quinoline Derivatives Using Copper Nanoparticles Grafted on Carbon Microspheres

Nitishkumar S. Kaminwar^a, Sunil U. Tekale^b, Rajkumar U. Pokalwar^c,
László Kótai^d, and Rajendra P. Pawar^e

^aDepartment of Chemistry, L.B.S. College, Dharmabad, India; ^bDepartment of Chemistry, Deogiri College, Aurangabad, India; ^cDepartment of Chemistry, Degloor College, Degloor, India; ^dResearch Centre for Natural Sciences, ELKH, Budapest, Hungary; ^eDepartment of Chemistry, Shiv Chhatrapati College, Cidco, Aurangabad, India

ABSTRACT

A rapid and efficient one-pot three-component synthesis of 2-amino-3-cyano-4-aryl-7-methyl-5-oxo-4,5-dihydro-pyrano-[3,2-c]pyran and 2-amino-3-cyano-4-aryl-6-methyl-5,6-dihydro-5-oxo-4H-pyrano-[3,2-c]quinoline derivatives catalyzed by copper nanoparticles grafted on carbon microspheres is developed. The use of heterogeneous catalyst, mild reaction conditions, and excellent yield of the corresponding products are the key features of the present protocol.

ARTICLE HISTORY

Received 18 July 2020
Accepted 26 February 2021

KEYWORDS

Cu-NP/C; pyranopyran;
pyranoquinoline

Introduction

Multicomponent reactions (MCRs) are the convergent reactions in which three or more reactants combine resulting in molecular complexity.¹⁻² MCRs are beneficial in terms of their atom economic, time saving, cost effective and environmentally benign nature.³ During the past decades, MCRs have attracted the attention of organic and medicinal chemists and constitute the efficient tools of green chemistry.⁴

4H-Pyran skeleton containing compounds constitute an important class of organic compounds exhibiting a wide range of pharmacological and biological properties.⁵ Pyran is one of the important structural motifs found widely in natural products, such as coumarins, benzopyrans, sugars, flavonoids, xanthenes, etc.⁶ Epicalyxins F and G, isolated from the seeds of *Alpinia blepharocalyx*, are the most potent members of this class that act as anticancer agents against human HT-1080 fibrosarcoma and murine 26-L5 carcinoma. Some biologically significant 4H-pyran derivatives are depicted in Figure 1. The synthesis of such compounds has attracted a strong interest owing to their biological potential.⁷ Various methods have been reported for the synthesis of these compounds and their analogues such as the use of 1H-imidazol-3-ium tricyanomethanide,⁸ iron oxide/phenyl sulfonic acid,⁹ PtCl₄,¹⁰ KF/Al₂O₃,¹¹ NH₄OAc,¹² ultrasonic irradiation,¹³ β-cyclodextrin,¹⁴ Fe₃O₄ nanoparticles,¹⁵ and ES/Cu(OH)₂ nanocomposite.¹⁶

Pyranoquinolines constitute an important group of heterocyclic compounds present in many natural products possessing different therapeutic activities.¹⁷ These compounds are biologically




Figure 1. Some representative examples of biologically potent pyrano-pyrans.


significant for exhibiting numerous activities such as antiallergic, antifungal, antibacterial, antipyretic, insecticidal, analgesic, antipyretic, cytotoxic, antiplatelet aggregation agents, and anticancer.¹⁸ Pyranoquinoline containing alkaloids such as fusaricide, melicobisquinolinone-B, zanthosimuline, and huajiaosimuline exhibit cytotoxicity against cancer cells as potential anticancer agents. Many methods involving one-pot three-component reaction of aldehydes, malononitrile, and 4-hydroxy-2-quinolone have been documented under the catalytic influence of Et_3N ,¹⁹ L-Proline,²⁰ DABCO,²¹ and CoFe_2O_4 ,²² Fe_3O_4 @GO-naphthalene- SO_3H ,²³ N-butylsulfonate-functionalized MWCNTs-D-NH₂,²⁴ Fe_3O_4 @DNH(CH₂)₄SO₃H,²⁵ SO₃H-functionalized nano-MGO-D-NH₂,²⁶ and [BMIM][HSO₄].²⁷ However, most of the methods are inconvenient due to certain drawbacks which include elaborate and tedious workup, use of toxic reagents, and large amounts of catalysts. Consequently, a new, eco-friendly, and low-cost protocol for the synthesis of pyranoquinoline derivatives is highly desirable.

Nano catalysts of copper are important nanoparticles.²⁸ Copper nanoparticle catalyzed reactions are advantageous over the conventional metal-catalyzed reactions in terms of low catalyst loading, high atom economy, better yields, inexpensive, shorter reaction times, and recyclability of the catalyst. Nanoparticles undergo self-aggregation and it is difficult to separate them from the reaction mixture. This limitation can be overcome by anchoring them on some supports. These developed catalytic metallic nanoparticles are inexpensive, non or minimally poisonous, highly active, stable, and easily separable from reaction mixture. Thus in continuation of our previous work on copper nanoparticles grafted on carbon microspheres,²⁹⁻³⁰ in the present work we report a simple and convenient method for the synthesis of 2-amino-3-cyano-4-aryl-7-methyl-5-oxo-4,5-dihydro-pyrano-[3,2-c]-pyran and 2-amino-3-cyano-4-aryl-6-methyl-4-hydro-pyrano-[3,2-c]-quinoline derivatives via one-pot three-component condensation of an aromatic aldehyde, malononitrile and the enolisable ketone, 4-hydroxy-6-methylpyran-2H-one/1-methyl-4-hydroxyquinoline-2H-one, under catalysis of copper nanoparticles grafted on carbon microspheres (Cu-NP/C) (Scheme 1).

Experimental section

General details

All the chemicals were Sigma Aldrich, Alpha Aeser or Spectrochem made and used without further purification. Thin Layer Chromatography (TLC) was monitored on Merck made silica gel



Scheme 1. One-pot synthesis of pyrano-pyran and pyrano-quinoline derivatives

60-F-254 aluminum plates. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker spectrometer using the specified solvents using tetramethylsilane as the internal standard. Mass spectrometric data were recorded by electron spray ionization (ES) technique on a Q-tof-micro quadrupole mass spectrometer (Micro mass).

General procedure for the synthesis of 6-amino-4-aryl-5-cyano-3-methyl-1-phenyl-1,4-dihydropyrano [2,3-c]pyrazoles


A mixture of aromatic aldehyde (1 mmol), malononitrile (1.1 mmol), 4-hydroxy-6-methyl-2-pyridone or 1-methyl-4-hydroxy-2-quinolone (1 mmol) was refluxed at 80°C with constant stirring in 1:1 water: ethanol (5 mL) in the presence of Cu-NP/C (10 wt%) for an appropriate time as mentioned in Table 2. The progress of the reaction was monitored by TLC using 5% methanol: dichloromethane as the mobile phase. After completion of the reaction, the reaction mixture was diluted with hot ethanol (10 mL) and filtered off to separate catalyst as the residue. The residue was further washed with hot ethanol (2×5 mL); the combined filtrates were concentrated on rotary evaporator to get the desired products which were of enough purity that required no purification.

Spectral data of the representative compounds is mentioned below:

2-Amino-3-cyano-4-(2-carboxyphenyl)-7-methyl-5-oxo-4,5-dihydro-5-oxopyrano[3,2-c]pyran, entry 9, (4h): MP: $238\text{--}240^\circ\text{C}$; IR (cm^{-1}) 752, 956, 1033, 1134, 1253, 1373, 1490, 1575, 1604, 1670, 2202, 2976, 3103, 3367, 3468; ^1H NMR (400 MHz, DMSO-d_6): δ ppm 2.20 (s, 3H, CH_3), 5.69 (s, 1H, CH, chiral), 6.25 (s, 1H, =CH), 7.09 (s, 2H, NH_2), 7.17-7.10 (d, 1H, Ar-H, $J=8.8$ Hz), 7.27-7.31 (t, 1H, Ar-H, $J=8.6$ Hz), 7.46-7.50 (t, 1H, ArH, $J=7.53$ Hz), 7.77-7.79 (d, 1H, ArH, $J=7.53$ Hz), 12.86 (s, 1H, COOH); HRMS (ES^+) m/z 325.07.

2-Amino-3-cyano-4-(4-cyanophenyl)-6-methyl-5,6-dihydro-5-oxo-4H-pyrano[3,2-c]quinoline, entry 14, (5b): MP: $278\text{--}280^\circ\text{C}$; IR (cm^{-1}) 754, 839, 1095, 1253, 1375, 1597, 1629, 1674, 2189, 2231, 3192, 3336, 3441; ^1H NMR (400 MHz, DMSO-d_6): δ ppm 3.53 (s, 3H, CH_3), 4.64 (s, 1H, CH chiral), 7.39 (s, 2H, NH_2), 7.42-7.44 (d, 3H, Ar-H, $J=8.52$), 7.57-7.59 (d, 1H, ArH, $J=8.00$), 7.71-7.76 (m, 3H, Ar-H), 8.02-8.04 (d, 1H, ArH, $J=7.2$); ^{13}C NMR (500 MHz, DMSO-d_6): δ ppm 29.1, 37.3, 38.8, 39.0, 56.6, 107.6, 109.4, 112.4, 114.6, 114.6, 119.2, 122.1, 128.5, 131.6, 132.2, 138.6, 149.7, 150.3, 158.7, 159.5; HRMS (ES^+) m/z 353.11 (M-1).

2-Amino-3-cyano-4-(4-(2-amino-3-cyano-5,6-dihydro-6-methyl-5-oxo-4H-pyrano[3,2-c]quinolin-4-yl)phenyl)-5,6-dihydro-6-methyl-5-oxo-4H-pyrano[3,2-c]quinoline (entry 15, 5c): MP: $279\text{--}282^\circ\text{C}$; IR (cm^{-1}) 752, 1095, 1290, 1377, 1462, 1627, 1672, 2194, 2229, 3190, 3327, 3371, 3431, 3587; ^1H NMR (400 MHz, DMSO-d_6): δ ppm 3.53 (s, 6H, 2CH_3), 4.47 (s, 1H, CH, chiral), 4.63 (s, 1H, CH, chiral), 7.10 (s, 1H, ArH), 7.23 (s, 1H, ArH), 7.38 (s, 4H, 2NH_2), 7.43-7.48 (d, 2H, ArH,



$J = 8.3$ Hz), 7.54-7.59 (m, 2H, Ar-H), 7.71-7.74 (m, 2H, ArH), 7.86-7.89 (m, 2H, ArH), 7.98-8.00 (m, 2H, ArH); ^{13}C NMR (500 MHz, DMSO- d_6): δ ppm 29.2, 29.2, 36.8, 37.5, 38.9, 40.0, 56.7, 57.8, 107.6, 108.9, 112.5, 113.2, 114.9, 119.3, 122.0, 114.1, 122.2, 127.3, 128.6, 129.8, 130.7, 131.7, 138.7, 150.1, 150.4, 151.1, 158.7, 159.7, 161.0; HRMS (ES^+) m/z 581.23.

Results and discussion

In the present work, we report an efficient and facile protocol for the synthesis of 2-amino-3-cyano-4-aryl-7-methyl-5-oxo-4,5-dihydro-pyrano[3,2-*c*]pyran and 2-amino-3-cyano-4-aryl-6-methyl-4-hydro-pyrano [3,2-*c*]quinoline derivatives by the one-pot three-component condensation from aromatic aldehydes, malononitrile, and 4-hydroxy 6-methyl-2-pyrone or 1-methyl-4-hydroxy-2-quinolone by using Cu NP/C as a reusable and safe catalyst in 1:1 water-ethanol mixture at 80°C.

The present work was carried by using Cu NP/C catalyst for the synthesis of pyrano-pyran and pyrano-quinoline derivatives. Initially, 3-bromo benzaldehyde (1 mmol), malononitrile (1.1 mmol), 4-hydroxy 6-methyl 2-pyrone (1 mmol) were selected for the optimization of reaction conditions at 80°C in a 5 mL water-ethanol (1:1). The effect of solvent and amount of catalyst was also studied (Table 1). A better yield was found using 10 wt% of the catalyst in 1:1 H₂O-EtOH (Table 1).

Thereafter, a series of reactions was carried out by using diversely substituted aldehydes under the optimized reaction conditions. All the aldehydes used underwent the three-component reaction with malononitrile and 4-hydroxy 6-methyl-2-pyrone to produce a library of 2-amino-3-cyano-4-aryl-7-methyl-5-oxo-4, 5-dihydro-pyrano [3,2-*c*]pyran derivatives in good to excellent yields. Similarly, 2-amino-3-cyano-4-aryl-6-methyl-4-hydro-pyrano [3,2-*c*]quinoline derivatives were synthesized by using 1-methyl 4-hydroxy 2-quinolone (Table 2).

Table 1. Optimization of the reaction conditions.

| Entry | Solvent | Catalyst (wt%) | Temperature (°C) | Time (min.) | Yield (%) |
|-------|---------------------------------|----------------|------------------|-------------|-----------|
| 1 | CH ₃ CN | 10 | 80 | 98 | 62 |
| 2 | CH ₂ Cl ₂ | 10 | 40 | 137 | 65 |
| 3 | H ₂ O | 05 | 80 | 114 | 67 |
| 4 | EtOH | 05 | 80 | 28 | 76 |
| 5 | 1:1 H ₂ O - EtOH | 05 | 80 | 25 | 80 |
| 6 | 1:1 H ₂ O - EtOH | 10 | 80 | 14 | 92 |

Table 2. Synthesis of 2-amino-3-cyano-4-aryl-7-methyl-5-oxo-4,5-dihydro-pyrano[3, 2-*c*] pyran and 2-amino-3-cyano-4-aryl-6-methyl-5,6-dihydro-5-oxo-4*H*-pyrano[3,2-*c*] quinoline derivatives catalyzed by copper nanoparticles grafted on carbon microspheres.

| Entry | Aldehyde ^a | Enolisable Ketone | Time (min.) | Product | Yield ^b (%) | Melting point (°C) |
|-------|--------------------------|-------------------|-------------|---------|------------------------|------------------------|
| 1 | Benzaldehyde | 3a | 20 | 4a | 86 | 249-251 ⁷ |
| 2 | 3-Bromo benzaldehyde | 3a | 18 | 4b | 88 | 258-260 ⁷ |
| 3 | 2-Fluorobenzaldehyde | 3a | 20 | 4c | 90 | 238-240 ⁷ |
| 4 | 2-Chloro benzaldehyde | 3a | 17 | 4d | 90 | 267-268 ¹¹ |
| 5 | 4-Cyano benzaldehyde | 3a | 15 | 4e | 88 | 230-232 ⁷ |
| 6 | 3-Nitro benzaldehyde | 3a | 15 | 4f | 93 | 234-235 ⁷ |
| 7 | 4-Bromo benzaldehyde | 3a | 18 | 4g | 88 | 214-215 ⁷ |
| 8 | 2-Carboxy benzaldehyde | 3a | 24 | 4h | 85 | 230-232 [Present work] |
| 9 | Thiophene-2-carbaldehyde | 3a | 30 | 4i | 84 | 242-243 ¹¹ |
| 10 | 4-Hydroxy benzaldehyde | 3b | 16 | 5a | 91 | 130-132 ¹⁸ |
| 11 | 4-Cyano benzaldehyde | 3b | 36 | 5b | 86 | 254-256 [Present work] |
| 12 | Terphthaldehyde | 3b | 24 | 5c | 88 | 279-282 [Present work] |
| 13 | Indole 3-carboxaldehyde | 3b | 28 | 5d | 85 | 192-194 ¹⁸ |
| 14 | Benzaldehyde | 3b | 20 | 5e | 91 | 153-154 ¹⁸ |
| 15 | 3-Nitro benzaldehyde | 3b | 22 | 5f | 90 | 190-192 ¹⁸ |

^aSubstituted aromatic aldehyde (1 mmol), malononitrile (1 mmol), enolisable ketone (1 mmol), and Cu NP/C (10 wt%), H₂O: EtOH (1:1) 5 mL, 80°C; ^bIsolated yield.




Table 3. Reusability of catalyst for the model reaction.

| Sr. No. | Run | Yield (%) |
|---------|-------|-----------|
| 1 | Fresh | 88 |
| 2 | I | 85 |
| 3 | II | 81 |
| 4 | III | 81 |





Figure 2. The plausible mechanism for the synthesis of pyrano-pyrans by using Cu-NP/C catalyst.

The reactions of aromatic aldehydes with electron-withdrawing substituents such as chloro and nitro groups proceeded at faster rates and gave better yields than those with electron-donating groups such as methoxy and methyl.

After completion of the reaction as monitored by TLC, the reaction mass was diluted with hot ethanol (3×5 mL) and the catalyst was separated by filtration. The combined filtrate was collected to get crude product which was purified by recrystallization in hot ethanol. Reusability is an important aspect of a heterogeneous catalyst. We studied the catalyst reusability in case of the model reaction of 3-bromobenzaldehyde, malononitrile, and 4-hydroxy-6-methyl-2-pyrone to form the corresponding pyrano-pyran (Entry 2, Table 3). The catalyst was found to work well in successive runs without an appreciable loss of its catalytic activity.



Scheme 2. The synthesis of catalyst.

Table 4. Comparison of literature methods for the synthesis of 1,4-dihydropyrano[2,3-c]pyran and 1,4-dihydropyrano[2,3-c]quinoline derivatives with the present protocol.

| Sr. No. | Conditions | Catalyst amount | Reference |
|---------|--|----------------------------|--------------|
| 1 | Magnetic iron oxide/phenylsulfonic acid, Distilled water, 50 °C, 10–45 min | 0.2 mol% | 8 |
| 2 | KF/Al ₂ O ₃ EtOH, R. T. 5–10 h | 50 mg / mmol of aldehyde | 10 |
| 3 | Ammonium acetate, Grinding, 10–18 min | 10 mol % | 11 |
| 4 | DABCO 80 °C, 20 min | 25 mol% | 21 |
| 5 | Nano Cobalt Ferrite EtOH, MW irradiation | 0.5 mg/10 mmol of aldehyde | 22 |
| 6 | Fe ₃ O ₄ @ NS-GO, water, R.T., 2–10 min | 0.01 g / mmol of aldehyde | 23 |
| 7 | MMWCNTs-D-(CH ₂) ₄ -SO ₃ H nanocomposite EtOH: H ₂ O, Reflux, 10–30 min | 0.04 g / mmol of aldehyde | 24 |
| 8 | [CH ₂ O(CH ₂) ₂ NH ₃] ₂ (CF ₃ COO) ₂ ionic liquid, solvent free, 80 °C, 12–30 min | 7 mol% | 31 |
| 9 | {[HM]C(CN) ₃ }, neat, 50 °C, 10–25 min | 2 mol% | 32 |
| 10 | Cu NP/C EtOH: H ₂ O (1:1), 80 °C, 15–30 min | 10 wt% | Present work |

Thus Cu-NP/C can be easily separated and reused several times successfully without a significant loss of activity. Another important aspect of the present work is the use of aqueous ethanol (1:1) as solvent for the reaction.

The plausible mechanism for the synthesis of pyrano-pyrans by using Cu-NP/C catalyst is depicted in Figure 2. Cu NPs are partially oxidized on surface by air. The Cu₂O generated on surface of the catalyst [(Cu₂O)Cu-NP/C] acts as a base to remove protons from malononitrile and the reacting 2H-pyrone, whereby it turns into (Cu⁺)Cu-NP/C and not into (Cu)Cu-NP/C. The Cu⁺ left on surface of the catalyst can act as Lewis acid to activate I and III toward the relevant nucleophilic attacks. (Cu⁺)Cu-NP/C is transformed into (Cu₂O)Cu-NP/C during protonation of intermediate anions. It involves formation of the intermediate (III) via condensation of an aldehyde with malononitrile. Simultaneously, 4-hydroxy-6-methyl-2-pyrone is transformed by the catalyst into the intermediate (IV), which undergoes a nucleophilic addition onto (III) in Michael manner to give the intermediate (V). The intermediate (V) undergoes cyclization followed by tautomerism to form the desired product (VII).

A similar mechanism can be proposed for the Cu-NP/C catalyzed synthesis of 2-amino-3-cyano-4-aryl-6-methyl-4-hydro-pyrano [3,2-c]quinolone by replacing the 4-hydroxy-6-methyl-2-pyrone with 1-methyl-4-hydroxy-2-quinolone.

The protocol for detailed synthesis and characterization of the catalyst is already reported in our previous work.^{29–30} It was prepared from a styrene-based divinylbenzene and acrylonitrile-modified and iminodiacetate (-CH₂N(CH₂COOH)₂) functionalized resin. The resin was treated with copper (II) sulfate solution and then the dried Cu-loaded resin was carbonized at 600 °C for 4 h in a high purity dry nitrogen stream (Scheme 2). A comparison of the present method with



course of the literature methods for the synthesis of these heterocyclic compounds is depicted in Table 9.

Conclusion

In summary, in the present work, we report an efficient and rapid method for one-pot three-component synthesis of 2-amino-3-cyano-4-aryl-7-methyl-5-oxo-4,5-dihydro-pyrano [3,2-c]pyrans and 2-amino-3-cyano-4-aryl-6-methyl-5,6-dihydro-5-oxo-4H-pyrano [3,2-c]quinolones catalyzed by Co-NP/C under heterogeneous conditions. The present method provides a valuable addition to the synthesis of these biologically significant heterocycles in terms of mild, environmentally benign reaction conditions and excellent yields of the products.

Acknowledgment

The authors are thankful to the Principal, L.B.S. College, Dharmabad, and the Principal, Devgiri College, Aurangabad.

Disclosure statement

No potential conflict of interest was reported by the authors.


ORCID

Sumit U. Telale <http://orcid.org/0000-0002-4370-1139>

Rajendra P. Pawar <http://orcid.org/0000-0002-1001-2000>

References

1. H. Alnoohad, M. Tefshakhoh, B. Malika, and J. P. Oush, "Acidic Ionic Liquid [H⁺NP][HSO₄]-Promoted One-Pot Synthesis of Dihydro-1H-Indeno[1,2-b]Pyridines and Polysubstituted Imidazoles," *Polycyclic Aromatic Compounds* 40, no. 5 (2020): 1465-500.
2. B. Malika, H. Nathayya, K. Tawher, H. Alnoohad, A. Amari, S. A. Houssem, and S. M. M. Noum, "Synthesis and Characterization of Nanosized Magnetic Co-Fe Mixed Oxide and Its Catalytic Behaviour towards One-Pot Synthesis of Polysubstituted Pyridine Derivatives," *Polycyclic Aromatic Compounds* 40, no. 3 (2020): 633-45.
3. H. Atharhat, A. Ecouvchaos, and B. Malika, "Green Synthesis of 3,4-Disubstituted Isoxazole 5(4H)-Ones in a Deep Eutectic Solvent," *Organic Preparations and Procedures International* 52, no. 6 (2020): 517-23.
4. A. Dongling, Wei Wang, and K. Wang, "Chemistry and Biology of Multicomponent Reaction," *Chemical Reviews* 112, no. 6 (2012): 5083-155.
5. Nguyen Dinh Thanh, Do Son Hai, Nguyen Thu Thu Ha, Do Tien Tung, Gao Thi Le, Hoang Thi Kim Van, Vu Ngoi Loan, Duong Ngoi Loan, and Le Huu Dang, "Synthesis, Biological Evaluation and Molecular Docking Study of 1,2,3,4-tetrahydro-5H-pyrano[2,3-d]pyrimidine as Potential Mycobacterium Tuberculosis Protein Tyrosine Phosphatase B Inhibitors," *Bioorganic & Medicinal Chemistry Letters* 29, no. 2 (2019): 164-71.
6. D. Kumar, P. Sharma, H. Singh, K. Nepali, G. K. Gupta, S. K. Jain, and F. Niu-Kang, "The Value of Pyrans as Anticancer Scaffolds in Medicinal Chemistry," *RSC Advances* 7, no. 59 (2017): 36977-99.
7. X. Fan, D. Feng, J. Qiu, X. Zhang, J. Wang, P. M. Lousseau, G. Andrei, R. Smoek, and E. De Clercq, "Practical and Efficient Synthesis of Pyrano[3,2-c]pyridane, Pyrano[4,3-b]pyran and their Hybrids with Nucleoside as Potential Antiviral and Anticancer Agents," *Bioorganic & Medicinal Chemistry Letters* 20, no. 1 (2009): 109-13.
8. Mohammad Ali Zaidiyev, Meyssam Yarec, Sacred Bagheri, Abbas Khoshnood, and Diego A. Alonso, "1H-Indenyl 5-Iodo-Triacetonemethanide ([HIM][ClCN₃]) as a Nanostructured Molten Salt Catalyst: Application to the Synthesis of Pyrano [4, 3-b] Pyrans," *Research on Chemical Intermediates* 43, no. 5 (2017): 3291-305.



9. D. Elhamifar, Z. Ramazani, M. Norouzi, and R. Mirbagheri, "Magnetic Iron Oxide/Phenylsulfonic Acid: A Novel, Efficient and Recoverable Nanocatalyst for Green Synthesis of Tetrahydrobenzo[b]Pyrans under Ultrasonic Conditions," *Journal of Colloid and Interface Science* 511, (2018): 392–401.
10. S. J. Pastine, S. W. Youn, and D. Sames, "Pt(IV)-Catalyzed Cyclization of Arene-Alkyne Substrates via Intramolecular Electrophilic Hydroarylation," *Organic Letters* 5, no. 7 (2003): 1055–8.
11. X. S. Wang, J. X. Zhou, Z. S. Zeng, Y. L. Li, D. Q. Shi, and S. J. Tua, "One-Pot Synthesis of Pyrano [3, 2-c] Pyran Derivatives Catalyzed by KF/Al_2O_3 ," *Arkivoc* 11, (2006) : 107–13.
12. D. Rajguru, B. S. Keshwal, and S. Jain, "Solvent-Free, Green and Efficient Synthesis of Pyrano [4, 3-b] Pyrans by Grinding and Their Biological Evaluation as Antitumor and Antioxidant Agents," *Medicinal Chemistry Research* 22, no. 12 (2013): 5934–9.
13. Ridaphun Nongrum, Geetmani Singh Nongthombam, Mattilang Kharkongor, Jims World Star Rani, Noimur Rahman, Chingrishon Kathing, Bekington Myrboh, and Rishanlang Nongkhlaw, "A Nano-Organocatalyzed Route towards the Efficient Synthesis of Benzo [b] Pyran Derivatives under Ultrasonic Irradiation," *RSC Advances* 6, no. 110 (2016): 108384–92.
14. E. A. Kataev, M. R. Reddy, G. N. Reddy, V. H. Reddy, C. S. Reddy, and B. V. Reddy, "Supramolecular Catalysis by β -Cyclodextrin for the Synthesis of Kojic Acid Derivatives in Water," *New Journal of Chemistry* 40, no. 2 (2016): 1693–7.
15. S. Asghari, and M. Mohammadnia, "Synthesis and Characterization of Pyridine-4-Carboxylic Acid-Functionalized Fe_3O_4 Nanoparticles as a Magnetic Catalyst for the Synthesis of Tetrahydrobenzo[b]Pyran Derivatives under Solvent-Free Conditions," *Inorganic and Nano-Metal Chemistry* 47, no. 7 (2017): 1004–11.
16. E. Mosaddegh, A. Hassankhani, and H. Karimi-Maleh, "Synthesis and Characterization of $ES/Cu(OH)_2$ Nanocomposite: A Novel and High Effective Catalyst in the Green Synthesis of Pyrano [4, 3-b]Pyrans," *Materials Science and Engineering: C* 46, (2015): 264–9.
17. J. P. Michael, "Quinoline, Quinazoline and Acridone Alkaloids," *Natural Product Reports* 20, no. 5 (2003): 476–93.
18. M. M. Abdou, "Chemistry of 4-Hydroxy-2 (1H)-Quinolone. Part 1: Synthesis and Reactions," *Arabian Journal of Chemistry* 10, (2017): S3324–S3337.
19. K. D. Upadhyay, N. M. Dodia, R. C. Khunt, R. S. Chaniara, and A. K. Shah, "Synthesis and Biological Screening of Pyrano [3, 2-c] Quinoline Analogs as anti-Inflammatory and Anticancer Agents," *ACS Medicinal Chemistry Letters* 9, no. 3 (2018): 283–8.
20. S. Zhu, J. Wang, Z. Xu, and J. Li, "An Efficient One-Pot Synthesis of Pyrano[3,2-c]Quinolin-2,5-Dione Derivatives Catalyzed by L-proline," *Molecules (Basel, Switzerland)* 17, no. 12 (2012): 13856–63.
21. R. Ghorbani-Vaghei, F. Rahmatpour, N. Sarmast, J. Mahmoudi, and A. Shahriari, "DABCO as a Green Catalyst for the Synthesis of Pyranoquinoline Derivatives," *Canadian Journal of Chemistry* 95, no. 5 (2017): 601–4.
22. S. Bandaru, R. K. Majji, S. Bassa, P. N. Chilla, R. Yellapragada, S. Vasamsetty, R. K. Jeldi, R. B. Korupolu, and P. D. Sanasi, "Magnetic Nano Cobalt Ferrite Catalyzed Synthesis of 4H-Pyrano [3, 2-h] Quinoline Derivatives under Microwave Irradiation," *Green and Sustainable Chemistry* 6, no. 2 (2016): 101–9.
23. M. Khaleghi-Abbasabadi, and D. Azarifar, "Magnetic Fe_3O_4 -Supported Sulfonic Acid-Functionalized Graphene Oxide ($Fe_3O_4@GO$ -naphthalene- SO_3H): a Novel and Recyclable Nanocatalyst for Green One-Pot Synthesis of 5-Oxo-Dihydropyrano[3,2-c]Chromenes and 2-Amino-3-Cyano-1,4,5,6-Tetrahydropyrano[3,2-c]Quinolin-5-Ones," *Research on Chemical Intermediates* 45, (2019): 2095–118.
24. F. Adibian, A. R. Pourali, B. Maleki, M. Baghayeri, and A. Amiri, "One-Pot Synthesis of Dihydro-1H-Indeno[1,2-b] Pyridines and Tetrahydrobenzo[b] Pyran Derivatives Using a New and Efficient Nanocomposite Catalyst Based on N-Butylsulfonate-Functionalized MMWCNTs-D- NH_2 ," *Polyhedron* 175, (2020): 114179.
25. Behrooz Maleki, Oliver Reiser, Ehsan Esmaeilnezhad, and Hyöung Jin Choi, " SO_3H -Dendrimer Functionalized Magnetic Nanoparticles ($Fe_3O_4@DNH(CH_2)_4SO_3H$): Synthesis, Characterization and Its Application as a Novel and Heterogeneous Catalyst for the One-Pot Synthesis of Polyfunctionalized Pyrans and Polyhydroquinolines," *Polyhedron* 162, (2019) : 129–41.
26. H. Alinezhad, M. Tarahomi, B. Maleki, and A. Amiri, " SO_3H -Functionalized Nano-MGO-D- NH_2 : Synthesis, Characterization and Application for One-Pot Synthesis of Pyrano[2,3-d]Pyrimidinone and Tetrahydrobenzo[b]Pyran Derivatives in Aqueous Media," *Applied Organometallic Chemistry* 33, no. 3 (2019): e4661.
27. K. Rad-Moghadam, S. C. Azimi, and E. Abbaspour-Gilandeh, "Synthesis of Novel Pyrano [3, 2-c] Quinoline-2, 5-Diones Using an Acidic Ionic Liquid Catalyst," *Tetrahedron Letters* 54, no. 35 (2013): 4633–6.
28. M. B. Gawande, A. Goswami, F.-X. Felpin, T. Asefa, X. Huang, R. Silva, X. Zou, R. Zboril, and R. S. Varma, "Cu and Cu-Based Nanoparticles: Synthesis and Applications in Catalysis," *Chemical Reviews* 116, no. 6 (2016): 3722–811.



29. T. Pasinszki, M. Krebsz, G. G. Lajgut, T. Koests, L. Kótal, S. Kauthale, S. Tekale, and R. Pawar, "Copper Nanoparticles Grafted on Carbon Microspheres as Novel Heterogeneous Catalysts and Their Application for the Reduction of Nitrophenol and One-Pot Multicomponent Synthesis of Hexahydroquinolines," *New Journal of Chemistry* 42, no. 2 (2018): 1092-8.
30. Y. W. More, S. U. Tekale, N. S. Kaminwar, L. Kótal, T. Pasinszki, P. S. Kendrekar, and R. P. Pawar, "Synthesis of 3,4-Dihydropyrano[c]chromenes Using Carbon Microsphere Supported Copper Nanoparticles (Cu-NP/C) Prepared from Loaded Cation Exchange Resin as a Catalyst," *Current Organic Synthesis* 16, no. 2 (2019): 288-93.
31. A. Zarei, M. Yarie, M. A. Zolfogol, and K. Niknam, "Synthesis of a Novel Bifunctional Oxyammonium-Based Ionic Liquid: Application for the Synthesis of Pyrano[4,3-b] Pyrans and Tetrahydrobenzo[b]Pyrans," *Journal of Chinese Chemical Society* 67, no. 6 (2020): 1120-31.
32. M. A. Zolfogol, M. Yarie, S. Baghery, A. Khoshnood, and D. A. Alonso, "1H-Imidazol-3-ium Tricyanomethanide $\{[HIM]C(CN)_3\}$ as a Nanostructured Molten Salt Catalyst: application to the Synthesis of Pyrano[4,3-b]Pyrans," *Research on Chemical Intermediate* 43, (2017): 3291-305.

Principal

A.V. Education Society
Degloor College Degloor