



Issue - XIV, Vol. V  
HI-TECH RESEARCH ANALYSIS

IMPACT FACTOR  
3.22

ISSN 2231- 6671  
Aug. 2017 To Jan. 2018



1

## IN-VITRO SCREENING OF SELECTED PLANT EXTRACTS ON SPORE GERMINATION OF HELIMINTHOSPORIUM SATIVUM

H. M. Lakde  
Dept. of Botany,  
Degloor College,  
Degloor, Dist. Nanded

Research Paper - Botany

### ABSTRACT

*In-vitro screening of ten selected plant extracts for antifungal activity against spore germination of Heliminthosporium Sativum, which causes leaf blight in Wheat. A study was undertaken to explore the effect of ten selected plant extract against spore germination of Heliminthosporium Sativum, which causes leaf blight in Wheat. In-Vitro study was down on effect of plant extracts on germination of spore. Inhibition of spore germination of Heliminthosporium Sativum seen at 10% concentration of leaf extract of some selected plants. Based on the study, it was concluded that some plant extract and plant products have potential for the control of phyto pathogenic fungi because fungal spores are main initial agent of diseases development.*

### Introduction :

The use of synthetic fungicides in plants disease management, no doute increases the yield of crop but has to hazardous effect on environment and human beings. The excessive and continuous uses of synthetic fungicides in disease management induce problems with development of resistance in pathogen against synthetic fungicides. The

  
**Dr. Anil Chidrawar**  
I/C Principal  
A.V. Education Society's  
Degloor College, Degloor Dist. Nanded



use of synthetic fungicides causes air, water and soil pollution, which is harmful to human being and useful micro product of soil including other non-pathogenic leaving organisms. By considering these drawbacks, the use of natural plant products in plant disease management came in to light from last century through out to world including India, the work on this aspect started in laboratory and at the field level. India the significant work done on use of neem, Karanja, Alive overa in plant disease management.

### MATERIAL AND METHODS

Plant materials used in testing are collected from local area and ground completely and filtered twice through Whatman filter paper. The plant product Neemarc also used for study. Plant extract and plant products were tested at 10 % concentration for antifungal activity.

Spore suspension of selected fungi was made by adding distilled water in to the petri-dishes which were kept for three weeks after inoculation at the temp. 27 $\pm$ 1 c. the concentration of selected spores was 10 mL measured by Hemacytometer (Hausser scientific, Hosham PA USA).

Plant extract of selected plant dropped on slides by micropipette with 20 $\mu$ L of spore suspension of selected fungi and 20 mL of each one of extracts. Each treatment contained three replication. The slides were put in to a plastic box with wet paper underneath in order to keep saturated humidity inside the box. The box was put into refrigerator at 4 c for 20 minutes and then into incubator for 24 hrs. At 27 $\pm$ 1 c. The germination rates of spores of selected fungi were count under the microscope. Result was mentioned in table I following table.

### RESULTS

The results of the present study regarding effect of ten selected plant extract and Neemarc as plant products deals with in vitro study of effect of plant extracts and plant products on germination of spores of *H. sativum*.

The effect of plant extract and plant products on germination of spores of selected fungi reveals that, the plant extract of *A. indica*, *A. vasica*, *A. sativum*, *C. gigantia*, *L. inermis*, *P. galbapara* shows strong inhibitory effect on germination of spores of *H. sativum*, which causes leaf blight in Wheat. The Neem seed oil and Neemarc, shows strong inhibitory



effect on germination of spores of *H. sativum* than the leaf extract of selected plants, at low concentration

### EFFECT OF TEN SELECTED PLANT EXTRACT ON SPORE GERMINATION OF HELIMINTHOSPORIUM SATIVUM

S. No.	Plant Species	Comparison	% inhibition
1	<i>Acalpha indica</i>	I	7.12
2	<i>Adhatoda vasica</i>	D	73.05
3	<i>Allium sativum</i>	F	70.00
4	<i>Azadirachta indica</i>	A	78.80
5	<i>Calotropis gigantea</i>	B	76.50
6	<i>Catharanthus roseus</i>	H	28.42
7	<i>Lowsonia inermis</i>	E	72.50
8	<i>Pangamia galbara</i>	C	73.40
9	<i>Vitax negundo</i>	G	65.80
10	<i>Lantana camera</i>	J	6.05
11	Dithane M-45 ( 0.25% )		95.80

### DISCUSSION AND CONCLUSION

In modern agriculture, for crop plant disease control, different types of synthetic toxic chemicals are used, which causes harmful hazards on ecosystem and on human life. To find alternatives to synthetic chemicals, from last two decades many investigators in the field of plant pathology try to investigate natural plant products, which are eco-friendly and low cost.

In the present investigation of Effect of plant extracts on spore germination of *H. sativum*, the in vitro study shows that plant extract of *A. indica*, *A. vasica*, *A. sativum*, *C. gigantea*, *L. inermis*, *P. galbabara* and *Neemarc* shows positive effect on spore germination of *Heliminthosporium Sativum*, which causes leaf blight diseases in Wheat.

Pandey V. N. And A. K. Srivastava (1995) recorded that, the growth of *Aspergillus*, *Penicillium* and *Paecilomyces* inhibited by oils obtained from *Cinnamomum cassia* and *Trachyspermum ammi*. Dr. v. Mariappan (1995) from Tamil Nadu Agricultural university Coimbatore, recorded positive efficiency of neem extract and Neem products on growth of *Pyricularia oryzae*, *Rhizoctonia solani*, *Heliminthosporium oryzae*,



*Cercospora persinata* and *Cercospora arachidicola*.

The same types of results are also confirmed by Herger et al (1988) i.e use of plant extract of Reynoutria and Olive oil for the control of powdery mildew of squash. The same results are also recorded by Haberic and Schlossor in 1993.

Clydette M Alsup (2004) done screening for active ingredient in plant extract that inhibit the growth of *A. tumefaciens* and concluded that the ingredient of *A. indica*, Jojoba oil, Cinnamon oil, Soyabin oil and Garlic can shows antimicrobial activity against *A. tumefaciens*, which causes crown galls in woody plants.

H. Ramezani, H. P. Singh, D. R. Batish, R. K. Kohli and J. S. Dargan in 2002 confirm the effect of volatile oils of *Eucalyptus citriodora* against two well-known rice pathogens, *Rhizoctonia solani* and *Helminthosporium oryzae*. They concluded that, complete inhibition of *R. solani* and *H. oryzae* was observed at 10 and 20 ppm concentration and from this study they also concluded that Eucalypt volatile oils have potential for the suppression of phytopathogenic fungi.

P. B. Bhende and V.V. Deshmukh (2003) recorded in vitro effect of leaf extract of Ashok and Kawach against *Alternaria alternata* which causes leaf spot diseases in sunflower and concluded that 70.98% of Kawach leaf extract inhibit growth of *Alternaria alternata* and 49.11% leaf extract of Ashok inhibit growth of *Alternaria alternata*.


Srinivas T, Rao K. C., Chattopadhyay C. (1997) confirm the effect of plant extracts on the management of blight of sunflower caused by *Alternaria alternata*, *Alternaria helianthi*. S

### References :-

- 1) Amadioha A. C. (2000): Controlling rice blast in vitro and in vivo with extracts of *Azadirachta indica*. *Crop prot.* 19: 287- 290
- 2) Appleton J. A., Tansey M. R. (1975)-Inhibition of growth of pathogenic fungi by garlic extracts. *Mycologia* 1975,67:882-885
- 3) Bisht G.S., Khulber R. D. (1995)- In vitro efficacy of leaf extract against leaf spots pathogeh of rice. *Indian phytopatho.* 48 (4) 480-482.
- 4) CAO Ke-Qiang, Ariena H C vans Bruggen (2001) -Inhibitory efficacy of several



- plant extracts and plant products on *Phytophthora infestans*. Journal of Agricultural University of Hebei 2001.
- 5) Chingduang, S., P. Siriacha and M. Saito. (1995). Effect of some medicinal plants and spices on growth of fungi. Pp. 143- 153.
  - 6) Dubey P.C., Sankaye K.S., and DESHMUKH M.B. (2003); Comparative study of *Allium sativum*, *A. indica* & *Acacia indica* for there antifungal activity. National conf. Proceeding p-71
  - 7) Dubey, N.K.; Tiwari, T.N.; Mandin, D.; Anrdriaboavunjy, H.; Chaumont, J.P. (2000); Antiungal properties of *Ocimum gratissimum* essential oil. (Ethyl cinnamate chemotype). *Fitoterapia* 71: 567-569
  - 8) Garg, k. L. (1995); Use of homeopathic drugs as antifungal agents for the protection of books and paper materials, pp.56-65.
  - 9) Gupta, R.P. and A. Singh (1983): effect of certain plant extract and chemicals on teliospore germination of *neovossia indica*. *Indian myco. Plant patho.* 13: 116-117
  - 10) Gawai D. U. and Usha K. Deshpande (2003): Effect of aqueous extract of *Allium sativum* and *Vitex negundo* on growth of *Deightoniella torulosum* with 0.2% Bavistin and Mancozeb. *Advances in Biotech. Research* 2003 p 98-101
  - 11) Khan, M.W., Alam, M.M., Khan, A.M. and Saxena, S.K. (1974) Effect of water soluble fractions of oil-cakes and bitter principles of neem on some fungi and nematodes. *Acta Bot. Indica* 2:120-128.
  - 12) Kishore N. Dubey N. K. (1998) Fungitoxicity of some higher plants against *Trichophyton rubrum* and *Epidermophyon floccosum*. *Nndian Journal of pharma. Sciences* 1998 Nav-Dec.50(6):323-5.
  - 13) Mariappan, V., Rajeswari, E. and Kamalakannan, A. (1995) Management of rice blast, *Pyricularia oryzae* by using neem (*Azadirachta indica*) amd other plant products. In: Mariappan, V. [Ed.] *Neem for the Management of Crop diseases*. Associated Publishing Co., New Delhi, India. Pp. 3-10.

  
**Dr. Anil Chidrawar**  
IC Principal  
A.V. Education Society's  
Degloor College, Degloor, Dist. Nanded